

AN EXTRAORDINARY SCATTERED BROAD EMISSION LINE IN A TYPE 2 QSO¹

GARY D. SCHMIDT², PAUL S. SMITH², CRAIG B. FOLTZ³, AND DEAN C. HINES²
 gschmidt, psmith, cfoltz, dhines@as.arizona.edu
To appear in the Astrophysical Journal Letters

ABSTRACT

An infrared-selected, narrow-line QSO has been found to exhibit an extraordinarily broad H α emission line in polarized light. Both the extreme width (35,000 km s⁻¹ full-width at zero intensity) and 3,000 km s⁻¹ redshift of the line centroid with respect to the systemic velocity suggest emission in a deep gravitational potential. An extremely red polarized continuum and partial scattering of the narrow lines at a position angle common to the broad-line emission imply extensive obscuration, with few unimpeded lines of sight to the nucleus.

Subject headings: quasars: emission lines — quasars: individual (2MASS J130005.3+163214) — galaxies: active — polarization

1. INTRODUCTION

If the Unified Scheme for AGN (e.g., Antonucci 1993) is to be equally successful for objects of high as well as low luminosity, there should exist a large number of QSOs whose optical/UV spectra are dominated by narrow emission lines (Type 2 QSOs or QSO-2s). Like their Seyfert counterparts, the broad emission-line regions of such objects are expected to lie within and behind a nominally toroidal obscuring structure, so that broad lines are most readily visible in light polarized by scattering off particles above and below the plane of obscuration. Though a few narrow-line objects with quasar luminosities have been found with broad permitted lines in infrared and/or polarized light (esp. *IRAS* QSOs: Hines et al. 1995, 1999; Young et al. 1996), it is clear that such highly-obscured nuclei are generally not present in classic lists of objects selected via UV excess.

Far more promising is the AGN catalog being compiled from the Two Micron All Sky Survey (2MASS; Cutri et al. 2001). The fundamental selection criterion of this sample is a color limit ($J - K_S > 2$) that weeds out stars while at the same time favoring extragalactic sources with substantial reddening toward their nuclei. This expectation has received strong support from the discovery of optical linear polarization in a sizeable fraction of 2MASS QSOs (in some cases $P \gtrsim 10\%$; Smith et al. 2002) as well as in the detection of large absorbing columns toward the associated nuclear X-ray sources (Wilkes et al. 2002).

The above studies found a surprisingly large number of 2MASS QSOs with the seemingly paradoxical characteristics of polarization/absorption coupled with prominent Type 1 (broad emission-line) total-flux spectra, properties that are not easily explained by the simplest models. Nevertheless, the narrow-line 2MASS AGN provide a number of high-luminosity analogues to Seyfert 2 galaxies where one might seek the broad, polarized emission lines that signal the presence of an accretion source powering the nucleus. This paper reports the first such discovery: a

narrow-line QSO with an extraordinarily broad scattered H α emission line and a polarized flux spectrum that may yield new information on the inner structure of AGN.

2. OBSERVATIONS

In survey observations, 2MASS J130005.3+163214 (hereafter 2M130005) showed a moderately bright and rather red ($B = 17.1$, $B - K_S = 5.24$) spectrum with a substantial starlight component and prominent narrow emission lines indicating a redshift of $z = 0.080$. The 2MASS photometry implies an absolute $2\mu\text{m}$ magnitude $M_{K_S} = -25.84$, well within the range of classical QSOs like the PG sample (Schmidt & Green 1983). The coordinates of 2M130005 do not correspond with a detection in the *IRAS* Point Source Catalog but the object is an 11.34 mJy FIRST radio source (Gregg et al. 1996) and it is known to be absorbed in X-rays (Wilkes et al. 2002). With a Galactic latitude of 79° , the R -band polarization of $P = 1.68\% \pm 0.14\%$ (Smith et al. 2002) is both intrinsic and tantalizing. Spectropolarimetry was obtained on three occasions between 2001 Apr. and 2002 Jul. using the instrument SPOL (Schmidt, Stockman & Smith 1992) attached to the 6.5 m MMT and 2.3 m Bok reflector of Steward Observatory. Due to the smaller collecting area of the Bok telescope and an inefficient initial primary mirror coating on the MMT, the earlier data are dwarfed in quality by the latest epoch from the MMT. All observations used a low-resolution grating providing one full order of spectral coverage, typically $\lambda\lambda 4250 - 8500$. An entrance slit of $1''.1$ (MMT; $3''.0$ for the Bok telescope) yielded a FWHM resolution of $\sim 17 \text{ \AA}$. At the alt-azimuth MMT, the instrument rotator was initially aligned with the parallactic angle and then set to track to a constant equatorial offset throughout the observational sequences. Standard calibration procedures included measurements of a completely polarized source and of interstellar polarization standard stars made with identical setups during the same telescope runs.

¹ A portion of the results presented here made use of the Multiple Mirror Telescope Observatory, a facility operated jointly by The University of Arizona and the Smithsonian Institution.

² Steward Observatory, The University of Arizona, Tucson AZ 85721

³ Multiple Mirror Telescope Observatory, The University of Arizona, Tucson, AZ 85721

Data from the individual epochs show qualitatively the same signatures and polarimetric differences that are entirely attributable to different aperture sizes, the latter resulting in variable amounts of dilution from the host galaxy of the QSO. The data were therefore coadded, weighting according to the quality of each epoch. Figure 1 shows the final results in four panels. A total of 4.1 hr of integration is represented in the figure, dominated by ~ 2 hr with high-reflectance MMT optics.

3. AN ENORMOUS, SCATTERED $H\alpha$ EMISSION LINE

The total flux spectrum in Figure 1 is that of an emission-line galaxy, with prominent [O III] $\lambda\lambda 4959, 5007$, [O I] $\lambda 6300$, [N II] $\lambda\lambda 6548, 6584$, [S II] $\lambda\lambda 6717, 6731$, and $H\beta$ superimposed on a predominantly stellar continuum. Narrow $H\alpha$ can also be distinguished in the [N II] blend. These lines have FWHM of ~ 1000 km s $^{-1}$, i.e., slightly resolved in our data. Measurements of the line strengths, including the results of deblending of the $H\alpha$ + [N II] $\lambda\lambda 6548, 6584$ complex, yield the following line ratios: $\log([O III]\lambda 5007/H\beta) = 1.03$, $\log([N II]\lambda 6584/H\alpha) = 0.14$, $\log([S II]\lambda 6717+6731/H\alpha) = -0.09$, and $\log([O I]\lambda 6300/H\alpha) = -0.62$. In the diagnostic diagrams of Veilleux & Osterbrock (1987), the gas responsible for the narrow-line emission is clearly being ionized by an AGN.

The outstanding discovery, however, is an extraordinarily broad emission hump in polarized flux, $q' \times F_\lambda$, centered at rest wavelength $\sim \lambda 6630$. Interpreting this as scattered broad $H\alpha$ emission, its width (18,000 km s $^{-1}$ FWHM, or $\sim 35,000$ km s $^{-1}$ full-width at zero intensity) is the largest yet found for scattered permitted lines, and equal to the broadest observed in *any* AGN: cf. the extreme broad-line radio galaxies 3C 332 (e.g., Eracleous et al. 1997) and 3C 382 (Osterbrock, Koski, & Phillips 1976). The broad $H\alpha$ line in 2M130005 is redshifted relative to the narrow lines by ~ 3000 km s $^{-1}$ and situated atop an extraordinarily red continuum. Polarized, narrow [O III] $\lambda 5007$, [N II]+ $H\alpha$ are also prominent. It is important to note that the [O I] and [S II] lines are absent in polarized light, even though the data quality is certainly adequate to detect the latter if it were of the same relative strength as the [O III] and [N II] lines. From the line profile comparison shown as an inset in Figure 1, it appears that [N II] $\lambda 6584$ is significantly weaker relative to narrow $H\alpha$ in the polarized light spectrum than in total light. Finally, despite all of the structure in polarized flux, the position angle rotates smoothly by less than 15° across the observed spectrum.

We have attempted to make a correction for the host galaxy's contribution to the total flux spectrum by scaling a spectrum of the E0 galaxy NGC 3379 (Kennicutt 1992) to the strength of the stellar absorption lines in 2M130005. Our best attempt at cancelling the G and Mg I b bands is depicted by the thin line denoted "AGN" in the bottom panel of Figure 1. The subtraction is not equally good for all features, but the resulting stellar fraction of 0.8 at $\lambda 6000$ is probably correct to within 10%. The Na I D feature could not be eliminated for any stellar fraction short of unity, and that choice oversubtracted the blue continuum. The fact that this feature is marginally apparent in polarized flux suggests that it may be intrinsic to the

scattered light⁴. It is important to note that the broad $H\alpha$ component can be discerned as a shallow hump underlying the narrow lines in the residual AGN spectrum. This is true for *any* reasonable stellar fraction. A dotted line indicates the underlying AGN continuum in Figure 1.

Comparison of the peak brightness of broad $H\alpha$ in polarized and total light, $\sim 1 \times 10^{-17}$ erg cm $^{-2}$ s $^{-1}$ Å $^{-1}$ vs. $4 - 5 \times 10^{-17}$ erg cm $^{-2}$ s $^{-1}$ Å $^{-1}$, implies a degree of polarization for the scattered light of $P \geq 20\%$, with the inequality resulting from the possibility that some of the broad line in the total flux spectrum is unscattered. (This result effectively rules out dichroic extinction – the interstellar mechanism – as the polarizing agent.) By contrast, the implied polarization of the starlight-subtracted AGN continuum is nearly wavelength-independent at $P \approx 10\%$ over 5000 – 8500 Å and the continuum-subtracted polarizations of narrow [O III] $\lambda 5007$ and the [N II]+ $H\alpha$ complex are $1.91\% \pm 0.20\%$ and $2.58\% \pm 0.33\%$, respectively. Comparing these values to the measurement for broad $H\alpha$, we infer that $\lesssim 10\%$ of the total light in the narrow lines is scattered. The discrepancy between broad-line and (galaxy-corrected) continuum polarization is reminiscent of the situation in some Seyfert galaxies (e.g., Kay 1994; Tran 1995), where dilution by hot stars may be an explanation (e.g., González Delgado et al. 1998).

4. DISCUSSION

Much has been made of the illumination of particles situated above and below the toroidal plane by the brilliant nucleus. For edge-on perspectives, the scattering of broad-line emission by these particles can provide an indirect view of the nucleus that we detect as polarized lines like the enormous $H\alpha$ feature in 2M130005. The radius of the inner edge of the torus is typically inferred by molecular/maser and hot dust emission at $R \sim 1$ pc. Further reprocessing occurs at mid- and far-infrared wavelengths in a possibly distinct structure of much larger size. Clouds in the inner torus are thought to be individually optically thick at most wavelengths (Krolik & Begelman 1988), but the filling factor may be low, so it is unclear what the overall vertical optical depth is to visible-light photons in the inner few hundred pc. If this region has significant transparency, some of the narrow-line emission from one lobe of a bipolar narrow line region will pass through the central plane to be scattered off particles in the other lobe. With narrow lines of [O III], [N II], and $H\alpha$ showing a reduced, but significant, fractional polarization at the same position angle as the scattered nuclear emission, it is useful to estimate the importance of this effect.

We write the fraction of the narrow-line emission emitted on one side of the nucleus that is subsequently scattered by the opposite lobe as

$$\frac{L_{\lambda,SC}}{L_{\lambda,EM}} \sim \frac{\Omega R e^{-\tau}}{4\pi} \quad (1)$$

where Ω is the solid angle of the scattering lobe as seen from the emitting lobe, τ is the effective vertical optical depth of the nuclear region, and R is the effective scattering efficiency of the lobe. We have assumed isotropic emission for this crude exercise. Assuming that the scattering process is also isotropic implies that the observed

⁴ See the similar feature in Mrk 231 (e.g., Smith et al. 1995).

fluxes are in the same ratio. At the end of §3, we found that the observed ratio was ~ 0.1 . The quantity R includes variables such as the optical depth in a lobe and the ratio of scattering to total extinction, and we have difficulty imagining an overall value $R > 0.5$. For this limit and τ small, we find $\Omega \gtrsim \pi$, i.e., a half-opening angle for a scattering lobe $\beta \gtrsim 60^\circ$.

While the opening angle is uncomfortably large in comparison to the observed angular extents of scattering and [O III]-emitting lobes in resolved AGN, an ionization gradient is expected to exist within each cloud, with high-ionization species like O^{++} predominately confined to the surface facing the nuclear engine. Hence, a scattering cloud may see brighter narrow lines than an edge-on observer. An internal ionization structure also suggests that the backscattering effect would be greater for high-ionization lines than for low-ionization lines, possibly explaining the absence of [O I] $\lambda\lambda 6300, 6363$ and [S II] $\lambda\lambda 6717, 6731$ in the polarized light of 2M130005 (see Figure 1, inset). Finally, because this explanation places no requirement on the radial location of the scattering particles, the broad-line emission and continuum could be scattered so close to the toroidal plane that the light has difficulty emerging over the “lip” of the dusty torus. The resulting extinction could account for the unusually red polarized continuum depicted in Figure 1.

A more attractive explanation for the observed polarization spectrum may be scattering in an extended narrow-line region that contains a global ionization gradient. Stratification of the narrow-line region has been established in radio-loud AGN (Hes, Barthel, & Fosbury 1993) and Seyfert galaxies (Filippenko & Halpern 1984), among others, and can naturally lead to polarization differences between species (e.g., Hines et al. 1999 and Tran, Cohen & Villar-Martin 2000 for the [O III] and [O II] lines in IRAS P09104+4109). Unfortunately, this mechanism provides no natural explanation for the extremely red continuum in polarized light. The 5000 – 8000 Å portion of the latter can be matched to the spectrum of a typical QSO using a Galactic reddening law with $A_V \sim 4$ mag. At shorter wavelengths the dereddened spectrum turns up too rapidly, suggesting the onset of a prominent 3000 Å bump. Of course, if dust grains are the scattering particles, this amount of extinction should be considered a lower limit, since the spectrum might be blued by the scatter-

ing. A somewhat smaller continuum suppression factor is inferred from a comparison of the measured equivalent width of [O III] $\lambda 5007$, 330 Å, with the average for low- z QSOs from Boroson & Green (1992), ~ 30 Å. However, this result is subject to possible extinction of the narrow-line emission. Heavy extinction of a scattered nuclear spectrum is apparently rare for AGN (see Young et al. 1996 for IRAS 2306+0505), and one is left with the impression that the nucleus of 2M130005 is obscured along most sightlines.

As noted above, the broad H α emission is redshifted by about 3000 km s $^{-1}$ with respect to the narrow lines and stellar features. This is substantially greater than the mean redward shift of H β with respect to the systemic velocity measured in the large samples of McIntosh et al. (1999) and Zamanov et al. (2002). In fact, the shift observed in 2M130005 is three times larger than the largest value reported in either study. This may indicate that scattering occurs in an outflowing medium (see also Hines et al. 1995). However, if we assume that the redshift has a gravitational origin, the ratio of the black hole mass to the characteristic radius of the broad line-emitting region can be estimated as $M_{BH}/R_{BLR} \simeq 10^6 M_\odot/\text{AU}$. It is interesting to note that this is similar to the result of an independent derivation, $M_{BH}/R_{BLR} \simeq 4 \times 10^5 M_\odot/\text{AU}$, made under the assumption that the broad-line clouds are in Keplerian orbits about the central black hole with a maximum orbital velocity equal to the half width at zero intensity of scattered H α ($\sim 18,000$ km s $^{-1}$). An estimate for the black hole mass in 2M130005 therefore lacks only a size for the broad-line region from, e.g., reverberation mapping (e.g., Kaspi et al. 2000) or the overall accretion luminosity. The latter will be possible using wide-band photometry with *SIRTF*. An X-ray spectrum will soon be obtained, and infrared spectroscopy will be essential for defining the intrinsic spectrum and obscuration of the nucleus of this new Type 2 QSO.

We are grateful to A. Marble for assistance with the observations, to J. McAfee for skillful operation of the new MMT, and to the referee A. Laor. This research was supported by *SIRTF*/MIPS and Science Working Group contracts 860785 and 959969 and NSF grants 97-30792 and 98-03072.

REFERENCES

- Antonucci, R. 1993, *ARA&A*, 31, 473
 Boroson, T.A., & Green, R.F. 1992, *ApJS*, 80, 109
 Cutri, R.M., Nelson, B.O., Kirkpatrick, J.D., Huchra, J.P. & Smith, P.S. 2001 in *ASP Conf. Ser.* 232, *The New Era of Wide Field Astronomy*, ed. R. Clowes, A. Adamson, & G. Bromage (San Francisco: ASP), 78
 Eracleous, M., Halpern, J.P., Gilbert, A.M., Newman, J.A., & Filippenko, A.V. 1997, *ApJ*, 490, 216
 Filippenko, A.V., & Halpern, J.P. 1984, *ApJ*, 285, 458
 Gonzalez Delgado, R.M., Heckman, T., Leitherer, C., Meurer, G., Krolik, J., Wilson, A.S., Kinney, A., & Koratkar, A. 1998, *ApJ*, 505, 174
 Gregg, M.D., Becker, R.H., White, R.L., Helfand, D.J., McMahon, R.G., & Hook I.M. 1996, *AJ*, 112, 407
 Hes, R., Barthel, P.D., & Fosbury, R.A.E. 1993, *Nature*, 362, 326
 Hines, D.C., Schmidt, G.D., Smith, P.S., Cutri, R., & Low, F.J. 1995, *ApJ*, 450, L1
 Hines, D.C., Schmidt, G.D., Wills, B.J., Smith, P.S., & Sowiński, L.G. 1999, *ApJ*, 512, 145
 Kaspi, S., Smith, P.S., Netzer, H., Maoz, D., Jannuzi, B.T., & Givon, U. 2000, *ApJ*, 533, 631
 Kay, L.E. 1994, *ApJ*, 430, 196
 Kennicutt, R.C. 1992, *ApJS*, 79, 255
 Krolik, J.H., & Begelman, M.C. 1988, *ApJ*, 329, 702
 McIntosh, D.H., Rix, H.-W., Rieke, M.J., & Foltz, C.B. 1999, *ApJ*, 517, L73
 Osterbrock, D.E., Koski, A.T., & Phillips, M.M. 1976, *ApJ*, 206, 898
 Schmidt, G.D., Stockman, H.S., & Smith, P.S. 1992, *ApJ*, 398, L57
 Schmidt, M., & Green, R.F. 1983, *ApJ*, 269, 352
 Smith, P.S., Schmidt, G.D., Allen, R., & Angel, J.R.P. 1995, *ApJ*, 444, 146
 Smith, P.S., Schmidt, G.D., Hines, D.C., Cutri, R.M., & Nelson, B.O. 2002, *ApJ*, 569, 23
 Tran, H.D. 1995, *ApJ*, 440, 597
 Tran, H.D., Cohen, M.H., & Villar-Martin, M. 2000, *AJ*, 120, 562
 Veilleux, S. & Osterbrock, D.E. 1987, *ApJS*, 63, 295
 Wilkes, B.J., Schmidt, G.D., Cutri, R., Ghosh, H., Hines, D., Nelson, B., & Smith, P.S. 2002, *ApJ*, 564, 65

- Young, S., Hough, J.H., Axon, D.J., Ward, M.J., & Bailey, J.A. 1996, MNRAS, 280, 291
- Zamanov, R., Marziani, P., Sulentic, J.W., Calvani, M., Dultzin-Hacyan, D., & Bachev, R. 2002, ApJ, 576, 9




FIG. 1.— Spectropolarimetry of 2M130005. From bottom to top: the observed total flux spectrum; the Stokes flux $q' \times F_{\lambda}$ for a frame aligned with the overall position angle of polarization; the rotated Stokes parameter q' (%); and the equatorial position angle θ . Also shown in the bottom panel is a scaled spectrum of the elliptical galaxy NGC 3379 used to subtract the stellar absorption features in 2M130005, as well as the resulting spectrum of the AGN itself. The inset compares the narrow H α spectra in total and polarized flux on arbitrary scales.